

Presence of β -Lactamase Gene TEM-1 DNA Sequence in Commercial *Taq* DNA Polymerase

The development and spread of expanded-spectrum β -lactamases (ESBLs) that cause resistance to β -lactam antibiotics has contributed to great concern worldwide. Most ESBLs are derived from TEM and SHV β -lactamases by point mutations within the *bla*_{TEM} and *bla*_{SHV} genes, giving rise to extended-spectrum drug resistance (2, 3). The standard method for determining the specific ESBL gene for the more than 90 TEM-type and the more than 25 SHV-type ESBLs is PCR followed by nucleotide sequencing (2).

We routinely applied the standard method to confirm that ESBL genes were present in clinical strains of *Enterobacteriaceae* (6). To avoid cross-contamination, we used separate rooms for sample preparation, PCR assembly, and agarose gel analysis. Recently, we noticed that the negative (water) controls used in PCR amplification for *bla*_{TEM} genes produced positive results; this did not occur when the *bla*_{SHV} gene was targeted. The PCR product was of the predicted size, and nucleotide sequencing revealed that it was the *bla*_{TEM-1} gene. After systematic analysis of pipette tips, microcentrifuge tubes, and reagents for PCR, we found that the *Taq* DNA polymerase was the source of contamination. As shown in Fig. 1, *Taq* DNA polymerase from manufacturer A produced a strong signal, but that from manufacturer B did not.

PCR is widely used to detect specific DNA sequences for purposes of microbial identification, clinical diagnosis, and basic research. Because the method is extremely sensitive, a small amount of contaminating DNA can be a serious problem. *Taq* DNA polymerase is often expressed as a recombinant protein in *Escherichia coli*. For studies involving gene cloning and protein expression in *E. coli*, the *bla*_{TEM-1} gene has been the most commonly used selective marker for expression vectors that are generally present in multiple copies (9). It is likely that during *Taq* DNA polymerase purification, the DNA containing the *bla*_{TEM-1} gene was not completely removed. This failure may not be a rare occurrence, but the contamination would be detected only if primers specific for *bla*_{TEM} gene were used.

Several reports have documented the presence of exogenous DNA in commercial *Taq* DNA polymerases (1, 4, 5, 7, 8, 10). Sources of the contaminating DNA have ranged from bacteria (1, 4) and phage-like DNA (7) to both prokaryotes and eukaryotes (10); in other studies, it was determined that the contaminating DNA was not from *E. coli* or *Thermus aquaticus* (5, 8). In all of these previous reports, PCR amplification was performed with universal primers for the highly-conserved 16S rRNA gene, whereas in the present study, amplification was done with primers targeting the *bla*_{TEM} gene. Nevertheless, investigators, especially those who work on TEM-type ESBLs, should be aware of the possibility that *Taq* DNA polymerase is contaminated with the *bla*_{TEM-1} gene.

This study was funded by grants MMH9331 and MMH9459 from Mackay Memorial Hospital, Taipei, Taiwan.




FIG. 1. PCR products amplified from negative (water) controls with primers targeting the *bla*_{TEM} gene. The product was run on a 1% agarose gel with 0.5× Tris-acetate-EDTA buffer. All reactions were performed in duplicate. Lane M, DNA size marker; lanes 1 and 2, PCR buffer from manufacturer B with *Taq* DNA polymerase from manufacturer B; lanes 3 and 4, PCR buffer from manufacturer B with *Taq* DNA polymerase from manufacturer A; lanes 5 and 6, PCR buffer from manufacturer A with *Taq* DNA polymerase from manufacturer B; lanes 7 and 8, PCR buffer from manufacturer A with *Taq* DNA polymerase from manufacturer A. The arrow indicates the amplified *bla*_{TEM-1} gene fragment.

REFERENCES

- Böttger, E. C. 1990. Frequent contamination of *Taq* polymerase with DNA. *Clin. Chem.* **36**:1258–1259.
- Bradford, P. A. 2001. Extended-spectrum beta-lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin. Microbiol. Rev.* **14**:933–951.
- Bush, K., G. A. Jacoby, and A. A. Medeiros. 1995. A functional classification scheme for beta-lactamases and its correlation with molecular structure. *Antimicrob. Agents Chemother.* **39**:1211–1233.
- Corless, C. E., M. Guiver, R. Borrow, V. Edwards-Jones, E. B. Kaczmarek, and A. J. Fox. 2000. Contamination and sensitivity issues with a real-time universal 16S rRNA PCR. *J. Clin. Microbiol.* **38**:1747–1752.
- Hughes, M. S., L. A. Beck, and R. A. Skuce. 1994. Identification and elimination of DNA sequences in *Taq* DNA polymerase. *J. Clin. Microbiol.* **32**:2007–2008.
- Liu, C. P., N. Y. Wang, C. M. Lee, L. C. Weng, H. K. Tseng, C. W. Liu, C. S. Chiang, and F. Y. Huang. 2004. Nosocomial and community-acquired *Enterobacter cloacae* bloodstream infection: risk factors for and prevalence of SHV-12 in multiresistant isolates in a medical center. *J. Hosp. Infect.* **58**: 63–77.
- Newsome, T., B. J. Li, N. Zou, and S. C. Lo. 2004. Presence of bacterial phage-like DNA sequences in commercial *Taq* DNA polymerase reagents. *J. Clin. Microbiol.* **42**:2264–2267.
- Rand, K. H., and H. Houck. 1990. *Taq* polymerase contains bacterial DNA of unknown origin. *Mol. Cell. Probes.* **4**:445–450.

9. **Sambrook, J., and D. W. Russell.** 2001. *Molecular cloning: a laboratory manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
10. **Schmidt, T. M., B. Pace, and N. R. Pace.** 1991. Detection of DNA contamination in *Taq* polymerase. *BioTechniques* **11**:176–177.

Chuen-Sheue Chiang

Chang-Pan Liu*

Li-Chuan Weng

Nai-Yu Wang

Microbiology Section

Department of Medical Research

Mackay Memorial Hospital

45 Min-Sheng Rd., Danshui 251

Taipei, Taiwan

Gwo-Jen Liaw

Department of Life Science

National Yang-Ming University

Taipei, Taiwan

*Phone: 886-2-28094661, ext. 2414

Fax: 886-2-28094679

E-mail: cpliu@ms1.mmh.org.tw